arXiv:1209.1297v1 [math.DS] 6 Sep 2012

# Invariant formulation of a variational problem

Imsatfia Moheddine

September 7, 2012

**Abstract**

In this paper we present a Hamiltonian formulation of multisymplectic type of an invariant variational problem on smooth submanifold of dimension $p$ in a smooth manifold of dimension $n$ with $p < n$.

## Introduction

Let $\mathcal{M}$ be a smooth manifold of dimension $n$ and $\mathcal{N}$ be an oriented smooth submanifold of dimension $p$ in $T_x\mathcal{M}$, for $x \in \mathcal{M}$. We study the case $n = 3$ and $p = 2$ in details and consider generalizations. Let $\Sigma$ be an oriented surface in $\mathcal{M}$, in each $M \in \Sigma$, denote by $P$ the tangent plane. We define the *Grassmannian* as the set of all these elements. We define the 2-form areolar action $\ell$ homogenous of degree zero in $y$ and we denote by $L$ a lagrangian homogenous of degree one in $y$ such that $\frac{dL}{dy} = \ell$. For some basis $(u_1, u_2)$ of $P$, and $(\theta^1, \theta^2)$ the dual basis, we define the action

$$\mathcal{L}(\Sigma) := \int_\Sigma L(x, u_1 \wedge u_2)\theta_1 \wedge \theta_2,$$

We use notions of mechanics and we construct the multisymplectic type of Hamiltonnian formulation. We prove that this description is conserved by graphs by comparison with the presentation of Cartan in [3]. Finally we study this for $n > 3$ and $p < n$.

## 1 The case presented by Cartan: a surface $\Sigma$ in a smooth manifold $\mathcal{M}$ of dimension 3

Let $\mathcal{M}$ be a smooth manifold of dimension 3. We define the Grassmannian bundle of planes in $T\mathcal{M}$ by:

$$Gr^2\mathcal{M} := \{(x, E) | x \in \mathcal{M}, \quad E \text{ oriented plane}^1 \text{ in } T_x\mathcal{M}\}$$

**Definition 1.1.** Assume that $\Lambda^2 T_x\mathcal{M} := (\Lambda^2 T_x^*\mathcal{M})^*$ and $\Lambda^2 T\mathcal{M} = \bigcup_{x \in \mathcal{M}} \Lambda^2 T_x\mathcal{M}$. Let $u_1, u_2 \in T_x\mathcal{M}$. We define $u_1 \wedge u_2 \in \Lambda^2 T_x\mathcal{M}$ by: $\forall \alpha, \beta \in T_x^*\mathcal{M}$

$$(\alpha \wedge \beta)(u_1 \wedge u_2) := \alpha \wedge \beta(u_1, u_2) = \alpha(u_1)\beta(u_2) - \alpha(u_2)\beta(u_1).$$

---

[1] $E$ is called “element” by Cartan.

We denote by $\pi_{Gr^2\mathcal{M}} : Gr^2\mathcal{M} \longrightarrow \mathcal{M}$ the canonical projection. Let $x = (x^1, x^2, x^3)$ be a local coordinates on $\mathcal{M}$ and $\left(\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}, \frac{\partial}{\partial x^3}\right)$ a basis of $T_x\mathcal{M}$, so a basis of $\Lambda^2 T_x\mathcal{M}$ is given by

$$\left(\frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2}, \frac{\partial}{\partial x^2} \wedge \frac{\partial}{\partial x^3}, \frac{\partial}{\partial x^3} \wedge \frac{\partial}{\partial x^1}\right).$$

Let $(u_1, u_2)$ be a direct basis of $E$, and $(x^1, x^2, x^3; y^{12}, y^{23}, y^{31})$ the coordinates on $\Lambda^2 T\mathcal{M}$ where $y^{12}, y^{23}$, $y^{31}$ are the coordinates representing $E$. Then, we have

$$u := u_1 \wedge u_2 = y^{12}\frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2} + y^{23}\frac{\partial}{\partial x^2} \wedge \frac{\partial}{\partial x^3} + y^{31}\frac{\partial}{\partial x^3} \wedge \frac{\partial}{\partial x^1}$$

**Remark 1.2.** We consider the 3-form $\omega := dx^1 \wedge dx^2 \wedge dx^3$. We have:

1. The kernel of $u \lrcorner \omega$ is the plane $E$.
2. $Gr^2_x\mathcal{M} \simeq (\Lambda^2 T_x\mathcal{M} \setminus \{0\})/\mathbb{R}^*+$.

We deduce that the homogeneous coordinates on the Grassmannian bundle over $\mathcal{M}$ are $(x; y) := (x^1, x^2, x^3; [y^{12} : y^{23} : y^{31}])$

$$Gr^2\mathcal{M} := \{(x; y) | x = (x^1, x^2, x^3) \in \mathcal{M} \text{ and } E = y = [y^{12} : y^{23} : y^{31}]\},$$

**Definition 1.3.** Let $\mathcal{M}$ be a differential manifold and $n \in \mathbb{N}$, we say that a $(n + 1)$-form $\Omega$ in $\mathcal{M}$ is *multisymplectic* if we have

1. $\Omega$ is nondegenerate, that is, $\forall M \in \mathcal{M}$, $\forall \xi \in T_M\mathcal{M}$, if $\xi \lrcorner \Omega_M = 0$, then $\xi = 0$).
2. $\Omega$ is closed, $d\Omega = 0$.

Any manifold $\mathcal{M}$ equiped with a multisymplectic form $\Omega$ is called a *multisymplectic manifold.*

**Definition 1.4.** ([11]): The $(n + 1)$-form $\Omega$ is called *pre-multisymplectic form* if it's closed. A differential manifold $\mathcal{M}$ equiped of $\Omega$ is called *pre-multisymplectique manifold.*

## 1.1 Lagrangian formulation

**Definition 1.5.** A 2-form *areolar action* is a 2-form $\ell$ on $Gr^2\mathcal{M}$, which can be written

$$\ell(x, y) := \ell_{12}(x, y)dx^1 \wedge dx^2 + \ell_{23}(x, y)dx^2 \wedge dx^3 + \ell_{31}(x, y)dx^3 \wedge dx^1.$$

where $\ell_{12}(x, y), \ell_{23}(x, y)$ and $\ell_{31}(x, y)$ are homogeneous of degree 0 in $y$ over $]0, +\infty[$.

**Definition 1.6.** We define a homogeneous Lagrangian $L : \Lambda^2 T\mathcal{M} \to \mathbb{R}$ as a continuous function of $\Lambda^2 T\mathcal{M}$, such that

1. $L$ is of class $\mathcal{C}^1$ over $\Lambda^2 T\mathcal{M} \setminus \sigma_0$ where $\sigma_0$ is the zero section.
2. $L$ is homogeneous of degree 1 in $y$ over $]0, +\infty[$.

**Remark 1.7.** If $u_1, u_2$ are two vectors of the element $E_x$ then $L(x, u_1 \wedge u_2)$ represent *"the infinitesimal area"* of the parallelepiped formed by $u_1$ and $u_2$.

Denote by $\Sigma$ an oriented surface in $\mathcal{M}$, let $(u_1, u_2)$ be a moving basis on $\Sigma$ and $(\theta_1, \theta_2)$ its dual basis, we define

$$\mathcal{L}(\Sigma) := \int_\Sigma L(x, u_1 \wedge u_2)\theta_1 \wedge \theta_2, \tag{1}$$

Thus, by Euler formula, we have

$$L(x,y) = \sum_{\alpha<\beta} \frac{\partial L}{\partial y^{\alpha\beta}}(x,y) y^{\alpha\beta} = \sum_{\alpha<\beta} \ell_{\alpha\beta}(x,y) y^{\alpha\beta}.$$

Where

$$\begin{cases} \frac{\partial L}{\partial y^{12}} = \ell_{12} \\ \frac{\partial L}{\partial y^{23}} = \ell_{23} \\ \frac{\partial L}{\partial y^{31}} = \ell_{31} \end{cases}$$

So

$$\ell(x,y) := \sum_{\alpha<\beta} \left( dx^\alpha \wedge dx^\beta \otimes \frac{\partial}{\partial y^{\alpha\beta}} \right) \lrcorner\ dL = \frac{\partial L}{\partial y^{12}} dx^1 \wedge dx^2 + \frac{\partial L}{\partial y^{23}} dx^2 \wedge dx^3 + \frac{\partial L}{\partial y^{31}} dx^3 \wedge dx^1.$$

We deduce that the inner product by $\sum_{\alpha<\beta} \left( dx^\alpha \wedge dx^\beta \otimes \frac{\partial}{\partial y^{\alpha\beta}} \right)$ is the canonical operator of $(\Lambda^2 T_x\mathcal{M})^* \to \Lambda^2 T_x^*\mathcal{M}$ and

$$\ell = \sum_{\alpha<\beta} \left( dx^\alpha \wedge dx^\beta \otimes \frac{\partial}{\partial y^{\alpha\beta}} \right) \lrcorner \quad dL$$

with condition $y^{\alpha\beta} + y^{\beta\alpha} = 0$.

**Definition 1.8.** A surface $\Sigma$ is called a critical point of $\mathcal{L}$, if and only if, for all compact $K \subset \mathcal{M}$, $K \cap \Sigma$ is a critical point of

$$\mathcal{L}_K(\Sigma) := \int_{\pi^{-1}_{Gr^2\mathcal{M}}(K) \cap T\Sigma} \ell$$

where, $T\Sigma = \{(x, T_x\Sigma) \in Gr^2\mathcal{M}; x \in \Sigma\}$.

The action (1) given,

$$\mathcal{L}(\Sigma) := \int_{T\Sigma} \ell = \int_{T\Sigma} \sum_{\alpha<\beta} \left( dx^\alpha \wedge dx^\beta \otimes \frac{\partial}{\partial y^{\alpha\beta}} \right) \lrcorner \quad dL.$$

## 1.2 Hamiltonian formulation (multisymplectic)

We write the *Legendre transform* function by

$$\begin{array}{ccc} \Lambda^2 T\mathcal{M} & \longrightarrow & \Lambda^2 T^*\mathcal{M} \\ (x,y) & \longmapsto & \left(x, \dfrac{\partial L}{\partial y}(x,y)\right), \end{array}$$

where $\frac{\partial L}{\partial y}(x,y) = \left(\frac{\partial L}{\partial y^{\alpha\beta}}(x,y)\right)_{\alpha<\beta}$. We consider a Hamiltonian function by

$$\mathcal{H}(x,P) = \sum_{\alpha<\beta} p_{\alpha\beta} y^{\alpha\beta} - L(x,y),$$

where $y$ is a solution of $\frac{\partial L}{\partial y}(x,y) = p_{\alpha\beta}$. Then, by Euler formula, we have

$$L(x,y) = \frac{\partial L}{\partial y^{\alpha\beta}}(x,y) y^{\alpha\beta} = \sum_{\alpha<\beta} p_{\alpha\beta} y^{\alpha\beta} \text{ where } p_{\alpha\beta} = \frac{\partial L}{\partial y^{\alpha\beta}}(x,y),$$

so the Hamiltonian $\mathcal{H}$ vanishes.

**Definition 1.9.** We say that the Lagrangian $L$ is nondegenerate if it's of class $\mathcal{C}^2$ over $\Lambda^2 T\mathcal{M} \setminus \{\sigma_0\}$ and $\frac{\partial^2 (L^2)}{\partial y^{\alpha\beta} \partial y^{\alpha'\beta'}} > 0$ out of $\sigma_0$. In particular $L^2$ is then a function stricly convexe.

**Lemma 1.10.** If the Lagrangian $L$ is nondegenerate, then,

$$\operatorname{rank}\left(\frac{\partial^2 L^2}{\partial y^{\alpha\beta} \partial y^{\alpha'\beta'}}\right) = 1 + \operatorname{rank}(\operatorname{Hess}(L)).$$

*Proof.* The Lagrangian $L$ is class $\mathcal{C}^1$ over $\Lambda^2 T_x\mathcal{M}$, and class $\mathcal{C}^\infty$ over $\Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\}$, thus $L^2$ is homogeneous of degree 2 and of class $\Lambda^2 T_x\mathcal{M}$ and of class $\mathcal{C}^\infty$ over $\Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\}$. For $L$ is nondegenerate, this leads to the convexity. It's easy to check that $\frac{\partial (L^2)}{\partial y^{\alpha\beta}}(0) = 0$. Note $T_y S_x = \ker\left(\frac{\partial L}{\partial y^{\alpha\beta}}(S_x)\right)$, if $(e_1, e_2)$ is a basis of $T_y S_x$ and $e_3 = y = y^{\alpha'\beta'} \frac{\partial}{\partial y^{\alpha\beta}}$, we denote by $(z^1, z^2, z^3)$ the coordinates in basis $(e_1, e_2, e_3)$, we have $\frac{\partial L}{\partial z^1} = \frac{\partial L}{\partial z^2} = 0$. More $d\left(\frac{\partial L}{\partial y^{\alpha\beta}}\right)(e_3) = y^{\alpha'\beta'} \frac{\partial}{\partial y^{\alpha'\beta'}} \left(\frac{\partial L}{\partial y^{\alpha\beta}}\right) = \frac{\partial^2 L}{\partial y^{\alpha\beta} \partial y^{\alpha'\beta'}} y^{\alpha'\beta'} = 0$ by homogeneity, which results in the coordinates $z^a$ by $\frac{\partial^2 L}{\partial z^3 \partial z^a} = 0 \;\; \forall a = 1,2,3$. Thus if we compute $\frac{\partial^2 (L^2)}{\partial y^{\alpha\beta} \partial y^{\alpha'\beta'}}$ out of $\sigma_0$. we find

$$\operatorname{rank}\left(\frac{\partial^2 L^2}{\partial z^a \partial z^b}\right)_{1 \leq a,b \leq 3} = 1 + \operatorname{rank} d\left(\frac{\partial L}{\partial y}\right) |_{TyS} \tag{2}$$

In particular $L$ is nondegenerate if and only if rank $d\left(\frac{\partial L}{\partial y}\right) |_{TyS_x} = 2$, so $\frac{\partial L}{\partial y} |_{S_x}$ is an immersion $S_x$. $\square$

**Proposition 1.11.** *We consider the Lagrangian* $L : \Lambda^2 T\mathcal{M} \to \mathbb{R}$ *is of class* $\mathcal{C}^k, \quad k \geq 2$. *We have the following statement:*
*If* $L$ *is nondegenerate, then, the image of the Legendre transform,* $\mathcal{N} := \{(x,p) \in \Lambda^2 T^*\mathcal{M} / p_{\alpha\beta} = \frac{\partial L}{\partial y^{\alpha\beta}}(x,y)$, *is a regular convexe hypersurface in* $\Lambda^2 T^*\mathcal{M}$.

*Proof.* For $x$ in $\mathcal{M}$, we define by

$$\mathcal{N}_x = \{p \in \Lambda^2 T_x^*\mathcal{M} / p_{\alpha\beta} = \frac{\partial L}{\partial y^{\alpha\beta}}(x,y), y \in \Lambda^2 T\mathcal{M}\}.$$

First we show that $\mathcal{N}_x$ is a surface in $\Lambda^2 T_x^*\mathcal{M}$, to simplify notations, we denote $L(x,y) = L(y)$. Note

$$S_x := \{y \in \Lambda^2 T_x\mathcal{M} | L^2(y) = 1\}$$

$S_x$ is a submanifold embedding, in fact, $\forall y \in S$ we have $y^{\alpha\beta}\frac{\partial L}{\partial y^{\alpha\beta}}(y) = L(y) = 1 \neq 0 \Rightarrow d(L^2)_y \neq 0$.
We know that $\frac{\partial L}{\partial y^{\alpha\beta}}(\lambda y) = \frac{\partial L}{\partial y^{\alpha\beta}}(y)$ thus the image of $\Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\}$ by $\frac{\partial L}{\partial y^{\alpha\beta}}$ is the same as that of $S_x$, because for every half-line, $D \subset \Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\}$, $D$ cut $S_x$ in a single point.
We show that $\frac{\partial L}{\partial y^{\alpha\beta}}(S)$ a manifold embedding and convex. For $L^2$ is a strictly convex function, thus the Legendre transform

$$\begin{array}{rcl} \dfrac{\partial(L^2)}{\partial y^{\alpha\beta}} : \Lambda^2 T_x\mathcal{M} & \longrightarrow & \Lambda^2 T_x^*\mathcal{M} \\ y & \longmapsto & \dfrac{\partial L^2}{\partial y^{\alpha\beta}}(y) \end{array}$$

is a global homeomorphism, for $L^2$ is of class $\mathcal{C}^\infty$ over $\Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\}$ thus $L^2 : \Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\} \longrightarrow \Lambda^2 T_x^*\mathcal{M} \setminus \{\sigma_0^*\}$ is a diffeomorphism, for $S$ is a submanifold embedding, thus $\frac{\partial L^2}{\partial y^{\alpha\beta}}(S_x)$ is so. Or $\frac{\partial(L^2)}{\partial y^{\alpha\beta}} |_{S_x} = 2L\frac{\partial L}{\partial y^{\alpha\beta}} |_{S_x} = 2\frac{\partial L}{\partial y^{\alpha\beta}} |_{S_x}$ (because $\forall y \in S_x$ we have $L(y) = 1$).On denote by

$$B_x := \{y \in \Lambda^2 T_x\mathcal{M} | L^2(y) \leq 1\}$$

Show that $\frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$ is a convex set. Let $P_0, P_0' \in \frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$. Show that

$$\forall t \in [0,1] \Rightarrow tP_0 + (1-t)P_0' \in \frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$$

We have $P_0, P_0' \in \frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$ so there exist $y_0, y_0' \in B_x$ such that $P_0 = \frac{\partial L}{\partial y^{\alpha\beta}}(y_0)$ and $P_0' = \frac{\partial L}{\partial y^{\alpha\beta}}(y_0')$ or $B_x$ is convex, thus $ty_0 + (1-t)y_0' \in B_x \Rightarrow L(ty_0 + (1-t)y_0') = 1$ and since $\frac{\partial L}{\partial y^{\alpha\beta}} : \Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\} \longrightarrow \Lambda^2 T_x^*\mathcal{M} \setminus \{\sigma_0^*\}$ is a diffeomorphism, then there exists an unique $P_0'' \in \Lambda^2 T_x^*\mathcal{M} \setminus \{\sigma_0^*\}$ (so there exists a unique $y_0'' \in \Lambda^2 T_x\mathcal{M} \setminus \{\sigma_0\})$ such that $t\frac{\partial L}{\partial y^{\alpha\beta}}(y_0) + (1-t)\frac{\partial L}{\partial y^{\alpha\beta}}(y_0') = P_0'' := \frac{\partial L}{\partial y^{\alpha\beta}}(y_0'')$ and since $L$ homogeneous of degree one, thus, $\forall y \in \Lambda^2 T_x\mathcal{M}\setminus\{\sigma_0\}$ it exists an unique $\lambda > 0$ such as $L(\lambda y) = 1$, so $\lambda = \frac{1}{L(y'')}$ which gives $\frac{1}{L(y'')}y_0'' \in B_x$. Then $t\frac{\partial L}{\partial y^{\alpha\beta}}(y_0) + (1-t)\frac{\partial L}{\partial y^{\alpha\beta}}(y_0') = \frac{\partial L}{\partial y^{\alpha\beta}}(y_0'') = \frac{\partial L}{\partial y^{\alpha\beta}}\left(\frac{1}{L(y'')}y_0''\right) \in \frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$, which shows the convexity of $\frac{\partial L}{\partial y^{\alpha\beta}}(B_x)$ thus $\partial B_x = S_x$ is a convex surface . ☐

**Remark 1.12.** If $\frac{\partial^2(L^2)}{\partial y^{\alpha\beta}\partial y^{\alpha'\beta'}} > 0$, then $\mathcal{N}_x$ is topologically a sphere in $\Lambda^2 T_x^*\mathcal{M}$, for all $x \in \mathcal{M}$.

## 1.3 Application

Let the 2-form $\theta = p_{12}dx^1 \wedge dx^2 + p_{23}dx^2 \wedge dx^3 + p_{31}dx^3 \wedge dx^1$ on $\Lambda^2 T^*\mathcal{M}$, then

$$d\theta = dp_{12} \wedge dx^1 \wedge dx^2 + dp_{23} \wedge dx^2 \wedge dx^3 + dp_{31} \wedge dx^3 \wedge dx^1,$$

We have $\theta$ is a multisymplectic form on $\Lambda^2 T^*\mathcal{M}$, and $d\theta|_{\mathcal{N}}$ is a pre-multisymplectic form.

**Theorem 1.13.** *We have $\Sigma \subset \mathcal{M}$ a critical point of $\int_{T\Sigma} L$ if and only if the image of $T\Sigma$ by the Legendre is a surface in $\mathcal{N}$ such that $\theta|_{\mathcal{N}} = 0$, otherwise $\frac{\partial L}{\partial y}(T\Sigma) \subset (\mathcal{N}; \theta \mid_{\mathcal{N}})$, if we denote by*

$$\beta = \sum_{\alpha<\gamma} p_{\alpha\gamma} dx^\alpha \wedge dx^\gamma = \theta \mid_{\mathcal{N}}$$

*then,*

$$\mathcal{L}(\Sigma) = \int_{\frac{\partial L}{\partial y}(T\Sigma)} \beta = \int_\Gamma \beta \tag{3}$$

*where $\Gamma := \frac{\partial L}{\partial y}(T\Sigma) \subset \mathcal{N}$.*

*Proof.* Easily we can see that

$$\left(\frac{\partial L}{\partial y}\right)^* \theta = \frac{\partial L}{\partial y^{12}}dx^1 \wedge dx^2 + \frac{\partial L}{\partial y^{23}}dx^2 \wedge dx^3 + \frac{\partial L}{\partial y^{31}}dx^3 \wedge dx^1 = \ell(x, y),$$

thus

$$\mathcal{L}(\Sigma) = \int_{T\Sigma} \ell = \int_{T\Sigma} \left(\frac{\partial L}{\partial y}\right)^* \theta,$$

since $\frac{\partial L}{\partial y} \mid_{T\Sigma}$ is an embedding, we get

$$\mathcal{L}(\Sigma) = \int_{T\Sigma} \left(\frac{\partial L}{\partial y}\right)^* \theta = \int_{\frac{\partial L}{\partial y}(T\Sigma)} \theta = \int_\Gamma \theta.$$

☐

### 1.4 Comparison with graphs

Let $\mathcal{M} = \mathbb{R}^2 \times \mathbb{R}$ and $\Sigma$ be the graph of a smooth function $f : \mathbb{R}^2 \to \mathbb{R}$. For $(x, y) = (x^1, x^2, f(x^1, x^2)) \in \Sigma$, the tangent bundle $T_{(x,y)}\Sigma$ is generated by

$$u_1 = \begin{pmatrix} 1 \\ 0 \\ \frac{\partial f}{\partial x^1} \end{pmatrix} \text{ and } u_2 = \begin{pmatrix} 0 \\ 1 \\ \frac{\partial f}{\partial x^2} \end{pmatrix}.$$

We have

$$\begin{aligned} u_1 \wedge u_2 &= \left( \frac{\partial}{\partial x^1} + \frac{\partial f}{\partial x^1} \frac{\partial}{\partial x^3} \right) \wedge \left( \frac{\partial}{\partial x^2} + \frac{\partial f}{\partial x^2} \frac{\partial}{\partial x^3} \right) \\ &= \frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2} - \frac{\partial f}{\partial x^1} \left( \frac{\partial}{\partial x^2} \wedge \frac{\partial}{\partial x^3} \right) - \frac{\partial f}{\partial x^2} \left( \frac{\partial}{\partial x^3} \wedge \frac{\partial}{\partial x^1} \right). \end{aligned}$$

Denote $y^{12} = 1$, $y^{23} = -\frac{\partial f}{\partial x^1}$ and $y^{31} = -\frac{\partial f}{\partial x^2}$. Let $F(x^1, x^2, f, df)$ be a Lagrangian density such that the action over an infinitesimal square $dx^1 \wedge dx^2$ is $F(x^1, x^2, f, df) dx^1 \wedge dx^2$. Then there exist a 2-form areolar action $\ell$, such that for any surface $\Sigma \subset \mathbb{R}^3$, we have

$$\int_\Omega F(x^1, x^2, f, df) dx^1 \wedge dx^2 = \mathcal{L}(\Sigma) = \int_{T\Sigma} \ell.$$

Indeed, if $(e_1, e_2)$ is a basis of $\mathbb{R}^2$ and $(u_1, u_2)$ is a basis of $T_x\Sigma$, then the infinitesimal action of $F$ on the parallelogram generated by $(e_1, e_2)$ is

$$F(x^1, x^2, f, df) dx^1 \wedge dx^2 (e_1, e_2) = F(x^1, x^2, f, df) dx^1 \wedge dx^2 (u_1, u_2) = F(x^1, x^2, f, df) y^{12}.$$

If we want

$$\sum_{1 \le \alpha < \beta \le 3} \ell_{\alpha\beta}(x, y) dx^\alpha \wedge dx^\beta (u_1, u_2) = \sum_{1 \le \alpha < \beta \le 3} \ell_{\alpha\beta}(x, y) y^{\alpha\beta} = L(x, y),$$

then, we have

$$L(x, y) = \sum_{1 \le \alpha < \beta \le 3} \frac{\partial L}{\partial y^{\alpha\beta}} y^{\alpha\beta} = \sum_{1 \le \alpha < \beta \le 3} \ell_{\alpha\beta} y^{\alpha\beta} = y^{12} F(x^1, x^2, f, df).$$

Since, $\frac{y^{23}}{y^{12}} = -\frac{\partial f}{\partial x^1}$, $\frac{y^{31}}{y^{12}} = -\frac{\partial f}{\partial x^2}$, then

$$L(x, y) = y^{12} F\left(x, f, -\frac{y^{23}}{y^{12}}, -\frac{y^{31}}{y^{12}}\right).$$

**Remark 1.14.** The generalization of the above formula to hypersurfaces in a smooth manifold $\mathcal{M}$ of dimension $n$ yields

$$L(x_1 ... x_n, [y]) = y^{1...n-1} F\left(x, f, -\frac{y^{2...n}}{y^{1...n-1}}, ..., -\frac{y^{n1...n-2}}{y^{1...n-1}}\right).$$

## 2 Formulation of the problem with a smooth submanifold of dimension $p$

Let $\mathcal{M}$ be a smooth manifold of dimension $n \in \mathbb{N}^*$, for all $x = (x_1, ..., x_n) \in \mathcal{M}$ and $p < n$, then a basis of $\Lambda^p T_x\mathcal{M}$ is

$$\left(\frac{\partial}{\partial x^{\imath_1}} \wedge ... \wedge \frac{\partial}{\partial x^{\imath_p}}\right)_{1 \leq \imath_1 < ... < \imath_p \leq n}.$$

Note $(x, y) = (x_1, ..., x_n, y^{\imath_1 \cdots \imath_p})_{1 \leq \imath_1 < ... < \imath_p \leq n}$ the coordinates on $\Lambda^p_D T_x\mathcal{M} \setminus \{\sigma_0\}$. Let $E$ an oriented element[2] in the tangent space $T_x\mathcal{M}$.
If $(u_1, ..., u_p)$ a direct basis of $E$ then we can assume that $u_1 \wedge ... \wedge u_p \in \Lambda^p T_x\mathcal{M} \setminus \{\sigma_0\}$ represents $E$, denote

$\Lambda^p_D T_x\mathcal{M} := \{u_1 \wedge ... \wedge u_p | (u_1, ..., u_p)$ direct basis of $E$, and oriented in $T_x\mathcal{M}\}$,

thus for $u \in \Lambda^p_D T_x\mathcal{M}$ we have

$$u := u_1 \wedge ... \wedge u_p = \sum_{1 \leq \imath_1 < ... < \imath_p \leq n} y^{\imath_1 \cdots \imath_p} \frac{\partial}{\partial x^{\imath_1}} \wedge ... \wedge \frac{\partial}{\partial x^{\imath_p}}.$$

**Definition 2.1.** For all $x \in \mathcal{M}$, we define the Grassmannian bundle by:

$$Gr^p_x\mathcal{M} := \{(x, E) | x \in \mathcal{M}, \quad E \text{ an oriented element in } T_x\mathcal{M}\}$$

**Remark 2.2.** We have

$$Gr^p_x\mathcal{M} \simeq (\Lambda^p_D T_x\mathcal{M} - \{\sigma_0\})/\mathbb{R}^*$$

Then the homogeneous coordinates on the Grassmannian bundle over $\mathcal{M}$ of dimension $p < n$, are $(x_1, ..., x_n; [y^{\imath_1 \cdots \imath_p}]_{1 \leq \imath_1 < ... < \imath_p \leq n})$ thus,

$$Gr^p\mathcal{M} := \{(x; [y]) := (x_1, ..., x_n; [y^{\imath_1 \cdots \imath_p}]_{1 \leq \imath_1 < ... < \imath_p \leq n}) | x = (x^1, ..., x^n) \in \mathcal{M}$$
$$E = (y) = [y^{\imath_1 \cdots \imath_p}]_{1 \leq \imath_1 < ... < \imath_p \leq n} \text{ oriented element in } T_x\mathcal{M}\}$$

with condition $y^{\imath_1 \cdots \imath_k \cdots \imath_{k'} \cdots \imath_p} + y^{\imath_1 \cdots \imath_{k'} \cdots \imath_k \cdots \imath_p} = 0$.

### 2.1 Lagrangian formulation

**Definition 2.3.** Let $E_x = (x, [y]) := (x_1, ..., x_n; [y^{\imath_1 \cdots \imath_p}]_{1 \leq \imath_1 < ... < \imath_p \leq n}) \in Gr^p\mathcal{M}$, we define a $p$-form areolar action $\ell$ on $Gr^p_x\mathcal{M}$ by

$$\ell(x, [y]) := \sum_{1 \leq \imath_1 < ... < \imath_p \leq n} \ell_{\imath_1 ... \imath_p}(x, y) dx_{\imath_1} \wedge ... \wedge dx_{\imath_p},$$

where $\ell_{\imath_1 ... \imath_p}$ are homogenous of degree 0 in $y$ over $]0, +\infty[$.

---

[2] The element is a vector subspace of $T_x\mathcal{M}$.

**Definition 2.4.** Let a Lagrangian $L : \Lambda^p_D T_x\mathcal{M} \to \mathbb{R}$ as a smooth function on $\Lambda^p_D T_x\mathcal{M}$ such as

1. $L$ is $\mathcal{C}^\infty$ over $L : \Lambda^p T_x\mathcal{M} \setminus \{\sigma_0\}$ where $\sigma_0$ is the vanish section.
2. $L$ is homogenous of degree 1 in $y$ on $]0, +\infty[$.

**Remark 2.5.** For all $x \in \mathcal{M}$, if $(u_1, ..., u_p)$ is a direct basis $E_x \in Gr^p_x\mathcal{M}$, then $L(x, u_1 \wedge ... \wedge u_p)$ represent *" the infinitesimal volume action"* formed by $p$ vectors $u_1, ..., u_p$.

Let $(\theta_1, ..., \theta_p)$ be the dual basis of $(u_1, ..., u_p)$, we define

$$\mathcal{L}(Gr^p_x\mathcal{M}) := \int_{TGr^p_x\mathcal{M}} L(x, u_1 \wedge ... \wedge u_p)\theta_1 \wedge ... \wedge \theta_p. \tag{4}$$

If we have $\ell_{\imath_1...\imath_p} = \frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}}$ with $1 \leq \imath_1 < ... < \imath_p \leq n$, then for the Euler formula we have $\sum_{1\leq\imath_1<...<\imath_p\leq n} \frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}} dx_{\imath_1} \wedge ... \wedge dx_{\imath_p} = \sum_{1\leq\imath_1<...<\imath_p\leq n} \ell_{\imath_1...\imath_p} dx_{\imath_1} \wedge ... \wedge dx_{\imath_p}$ thus

$$\begin{aligned} \ell(x; [y]) &= \sum_{1\leq\imath_1<...<\imath_p\leq n} \left(dx_{\imath_1} \wedge ... \wedge dx_{\imath_p} \otimes \frac{\partial}{\partial y^{\imath_1\cdots\imath_p}}\right) \lrcorner \; dL \\ &= \sum_{1\leq\imath_1<...<\imath_p\leq n} \frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}} dx_{\imath_1} \wedge ... \wedge dx_{\imath_p}. \end{aligned}$$

Then $\sum_{1\leq\imath_1<...<\imath_p\leq n} \left(dx_{\imath_1} \wedge ... \wedge dx_{\imath_p} \otimes \frac{\partial}{\partial y^{\imath_1\cdots\imath_p}}\right)$ is the canonical operator $(\Lambda^p_x T\mathcal{M})^* \longrightarrow \Lambda^p T^*_x\mathcal{M}$, thus we write

$$\ell(x; [y]) = \sum_{1\leq\imath_1<...<\imath_p\leq n} \left(dx_{\imath_1} \wedge ... \wedge dx_{\imath_p} \otimes \frac{\partial}{\partial y^{\imath_1\cdots\imath_p}}\right) \lrcorner \; dL$$

Let $L$ be a Lagrangian, thus we can write the action (4),

$$\mathcal{L}(Gr^p_x\mathcal{M}) := \int_{TGr^p_x\mathcal{M}} \ell = \int_{TGr^p_x\mathcal{M}} \sum_{1\leq\imath_1<...<\imath_p\leq n} \left(dx_{\imath_1} \wedge ... \wedge dx_{\imath_p} \otimes \frac{\partial}{\partial y^{\imath_1\cdots\imath_p}}\right) \lrcorner \; dL.$$

**Definition 2.6.** A critical point of $\mathcal{L}$ if and only if, for all compact $K \subset \mathcal{M}$, $K \cap Gr^p_x\mathcal{M}$ is a critical point of $\mathcal{L}_K(Gr^p_x\mathcal{M}) := \int_{\pi^{-1}_{TGr^p\mathcal{M}}(K)\cap TGr^p_x\mathcal{M}} \ell$, where $\pi_{Gr^p\mathcal{M}} : Gr^p\mathcal{M} \longrightarrow \mathcal{M}$ is the canonical projection.

## 2.2 Legendre transform-Hamiltonian formulation (multisymplectic)

We consider the Hamiltonian function

$$\mathcal{H}(x,P) = \sum_{1\leq \imath_1<\ldots<\imath_p\leq n} p_{\imath_1\ldots\imath_p} y^{\imath_1\cdots\imath_p} - L(x,y)$$

where $\frac{\partial L}{\partial y}(x,[y]) = \left(\frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}}(x,y)\right)_{\imath_1<\ldots<\imath_p}$ is the Legendre transform and $y$ is a solution of $\frac{\partial L}{\partial y}(x,[y]) = p_{\imath_1\ldots\imath_p}(x,[y])$.
Since $L$ is homogenous of degree 1 in $y$ on $]0,+\infty[$, then by formula Euler we have

$$L(x,[y]) = \frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}}(x,y) y^{\imath_1\cdots\imath_p} = \sum_{1\leq \imath_1<\ldots<\imath_p\leq n} p_{\imath_1\ldots\imath_p} y^{\imath_1\cdots\imath_p}$$

we conclude that the Hamiltonian $\mathcal{H}$ vanishes.

**Proposition 2.7.** *We consider $L : \Lambda^p_D T\mathcal{M} \to \mathbb{R}$, a Lagrangian nondegenerate, then the image of Legendre transform $\mathcal{I} := \{(x,p) \in \Lambda^p_D T^*\mathcal{M} / p_{\imath_1\ldots\imath_p}(x,[y]) = \frac{\partial L}{\partial y^{\imath_1\cdots\imath_p}}(x,[y]), y \in \Lambda^p_D T\mathcal{M}\}$ is a regular hypersurface and convex in $\Lambda^p_D T^*\mathcal{M}$.*

*Proof.* Same proof of (1.11). ☐

On $\Lambda^p T^*\mathcal{M}$, we define the $p$-form $\theta$ by

$$\theta = \sum_{1\leq \imath_1<\ldots<\imath_p\leq n} p_{\imath_1\ldots\imath_p} dx_{\imath_1} \wedge \ldots \wedge dx_{\imath_p},$$

**Remark 2.8.** $\theta$ is a multisymplectic form on $\Lambda^P_D T^*\mathcal{M}$ and $d\theta|_{\mathcal{I}}$ is a pre-multisymplectic form.

**Theorem 2.9.** *A critical point of $\int_{TGr^p_x\mathcal{M}} L$ if and only if the image by the Legendre of $TGr^p_x\mathcal{M}$ is a vector subspace in $\mathcal{M}$, such that $\theta \mid_{\mathcal{I}}$ is vanishes. If*

$$\beta = \theta \mid_{\mathcal{I}} \text{ and } \Gamma := \frac{\partial L}{\partial y}(TGr^p_x\mathcal{M}) \subset \mathcal{I}$$

*then*

$$\mathcal{L}(Gr^p_x\mathcal{M}) = \int_{\frac{\partial L}{\partial y}(TGr^p_x\mathcal{M})} \beta = \int_{\Gamma} \beta. \tag{5}$$

*Proof.* See proof of (1.13). ☐

### 2.3 Comparison with graphs

We will take $\mathcal{M} = \mathbb{R}^p \times \mathbb{R}^{n-p}$ and $\Gamma$ the graph of smooth function $f : \mathbb{R}^P \to \mathbb{R}^{n-p}$. Denote $\left(q_\imath^\jmath := \frac{\partial f_\jmath}{\partial x_\imath}\right)_{\substack{1\leq\imath\leq p \\ 1\leq\jmath\leq n-p}}$, in a neighberhood of a point $x = (x_1, ..., x_p, f_1, ..., f_{n-p}) \in \Gamma$ the tangent bundle $T_x\Gamma$ is generated by

$$u^1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ \frac{\partial f_1}{\partial x_1} \\ \frac{\partial f_2}{\partial x_1} \\ \vdots \\ \frac{\partial f_{n-p}}{\partial x_1} \end{pmatrix}, u^2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_2} \\ \vdots \\ \frac{\partial f_{n-p}}{\partial x_2} \end{pmatrix}, ..., u^p = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ \frac{\partial f_1}{\partial x_p} \\ \frac{\partial f_2}{\partial x_p} \\ \vdots \\ \frac{\partial f_{n-p}}{\partial x_p} \end{pmatrix}$$

so that

$$u_1 \wedge u_2 \wedge ... \wedge u_p = \left(\frac{\partial}{\partial x_1} + \sum_{\jmath=p+1}^{n} \frac{\partial f_1}{\partial x_{\jmath-p}} \frac{\partial}{\partial x_\jmath}\right) \wedge \left(\frac{\partial}{\partial x_2} + \sum_{\jmath=p+1}^{n} \frac{\partial f_2}{\partial x_{\jmath-p}} \frac{\partial}{\partial x_\jmath}\right) \wedge ... \wedge$$

$$\left(\frac{\partial}{\partial x_p} + \sum_{\jmath=p+1}^{n} \frac{\partial f_p}{\partial x_{\jmath-p}} \frac{\partial}{\partial x_\jmath}\right) = \sum_{1\leq\imath_1<...<\imath_p\leq n} c^{\imath_1\cdots\imath_p} \frac{\partial}{\partial x_{\imath_1}} \wedge ... \wedge \frac{\partial}{\partial x_{\imath_p}}$$

with $c^{1..p} = 1$ and $c^{\imath_1\cdots\imath_p}$ (where $\imath_1...\imath_p \neq 1...p$) are functions which can be computed, and which depends on $\left(\frac{\partial f_\jmath}{\partial x_\imath}\right)_{\substack{1\leq\imath\leq p \\ 1\leq\jmath\leq n-p}}$. In particular, we can show that

$$\begin{cases} c^{12..p} = 1 \\ \frac{c^{1...\imath-1\imath+1...pp+\jmath}}{c^{123...p}} = \frac{\partial f_\jmath}{\partial x_\imath} \text{ for } 1 \leq \imath \leq p, \quad 1 \leq \jmath \leq n-p \end{cases}$$

We suppose $y^{\imath_1\cdots\imath_p} = c^{\imath_1\cdots\imath_p}$, $(e_1, ..., e_p, e_{p+1}, ..., e_n)$ is a basis of $\mathbb{R}^n \times \mathbb{R}^{n-p}$ and $(u_1, ..., u_p)$ a basis of $T_x\Gamma$.
If $F(x_1, ..., x_p, f_1, ..., f_{n-p}, \nabla f)$ is a Lagrangian density, then its infinitesimal action over the parallelepiped generated by $(e_1, ..., e_p)$ is

$$F(x_1, ..., x_p, f_1, ..., f_{n-p}, \nabla f) dx_1 \wedge ... \wedge dx_p(e_1, ..., e_p) =$$

$$F(x_1, ..., x_p, f_1, ..., f_{n-p}, \nabla f) dx_1 \wedge ... \wedge dx_p(u_1, ..., u_p) = F(x_1, ..., x_p, f_1, ..., f_{n-p}, \nabla f) y^{1...p}$$

But we ask that this precedent egality is equivalent to

$$\sum_{\substack{1\leq\imath\leq p \\ 1\leq\jmath\leq n-p}} \ell_{1...\imath-1\imath+1...pp+\jmath}(x, [y]) dx_1 \wedge ... \wedge dx_{\imath-1} \wedge dx_{\imath+1} ... \wedge dx_p \wedge dx_{p+\jmath}(u_1, ..., u_p)$$

$$\sum_{\substack{1\leq\imath\leq p\\1\leq\jmath\leq n-p}} \ell_{1\ldots\imath-1\imath+1\ldots pp+\jmath}(x,[y])y^{1\ldots\imath-1\imath+1\ldots pp+\jmath} = L(x,[y^{1\ldots\imath-1\imath+1\ldots pp+\jmath}])$$

then

$$L(x;[y]) = \sum_{\substack{1\leq\imath\leq p\\1\leq\jmath\leq n-p}} \frac{\partial L}{\partial y^{1\ldots\imath-1\imath+1\ldots pp+\jmath}} y^{1\ldots\imath-1\imath+1\ldots pp+\jmath}$$

$$= \sum_{\substack{1\leq\imath\leq p\\1\leq\jmath\leq n-p}} \ell_{1\ldots\imath-1\imath+1\ldots pp+\jmath}(x;[y])y^{1\ldots\imath-1\imath+1\ldots pp+\jmath} = y^{1\ldots p}F(x^1,\ldots,x^p,f_1,\ldots,f_{n-p},\nabla f)$$

thus

$$L(x,[y]) = y^{1\ldots p}F\left(x^1,\ldots,x^p,f_1,\ldots,f_{n-p},\left(\frac{y^{1\ldots\imath-1\imath+1\ldots pp+\jmath}}{y^{123\ldots p}}\right)_{\substack{1\leq\imath\leq p\\1\leq\jmath\leq n-p}}\right).$$

# References

[1] Thierry Aubin. *A course in differential geometry*, volume 27 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.

[2] Robert L. Bryant. Some remarks on finsler manifolds with constant flag curvature. *Houston J. Math.*, 28(2):221–262, 2002.

[3] E. Cartan. *Les espaces métriques fondés sur la notion d'aire.* Hermann. 6, rue de la Sorbonne, Paris, 1933.

[4] E. Cartan. *Les espaces de Finsler.* Hermann. 6, rue de la Sorbonne, Paris, 1934.

[5] Élie Cartan. *Les systèmes différentiels extérieurs et leurs applications géométriques.* Actualités Sci. Ind., no. 994. Hermann et Cie., Paris, 1945.

[6] Matias Dahl. An brief introduction to finsler geometry. 2006.

[7] Frédéric Hélein et Joseph Kouneiher. Covariant hamiltonian formalism for the calculus of variations with several variables: Lepage-dedecker versus de donder-weyl. *Adv. Theor. Math. Phys*, 8:565–601, 2004.

[8] Dikanaina Harrivel. Hamiltonian multisymplectic formalism and monge-ampere equation. 2006.

[9] Frédéric Hélein. Multisymplectic formalism and the covariant phase space. In *Variational problems in differential geometry*, volume 394 of *London Math. Soc. Lecture Note Ser.*, pages 94–126. Cambridge Univ. Press, Cambridge, 2012.

[10] Frédéric Hélein and Joseph Kouneiher. The notion of observable in the covariant Hamiltonian formalism for the calculus of variations with several variables. *Adv. Theor. Math. Phys.*, 8(4):735–777, 2004.

[11] Frédéric Hélein. Multisymplectic formalism and the covariant phase space. 2010.

[12] Thomas A. Ivey and J. M. Landsberg. *Cartan for beginners: differential geometry via moving frames and exterior differential systems*, volume 61 of *Graduate Studies in Mathematics.* American Mathematical Society, Providence, RI, 2003.

[13] Michael Spivak. *A comprehensive introduction to differential geometry. Vol. (I-II-III-IV-V).* Publish or Perish Inc., Wilmington, Del., second edition, 1979.